

Flexible Generation of E-Learning Exams in R: Moodle Quizzes, OLAT Assessments, and Beyond

Achim Zeileis, Nikolaus Umlauf, Friedrich Leisch

<http://eeecon.uibk.ac.at/~zeileis/>

Overview

- Motivation and challenges
- R package **exams**
- Exercises
- Exams
 - Combination of exercises
 - PDF output
 - HTML output
 - XML for **Moodle** or **OLAT**
- Discussion

Motivation and challenges

Motivation:

- Introductory statistics and mathematics courses for business and economics students at WU Wien and Universität Innsbruck.
- Courses are attended by more than 1,000 students per semester.
- Several lecturers teach lectures and tutorials in parallel.
- Need for integrated teaching materials: Presentation slides, collections of exercises, exams, etc.

Challenges:

- *Scalable exams*: Automatic generation of a large number of different exams, both written and online.
- *Associated self-study materials*: Collections of exercises and solutions from the same pool of examples.
- *Joint development*: Development and maintenance of a large pool of exercises in a multi-author and cross-platform setting.

R package exams

Tools chosen: R (for random data generation and computations) and \LaTeX (for mathematical notation) \Rightarrow Sweave.

Design principles of package exams:

- Each exercise template (also called “exercise” for short) is a single Sweave file (`.Rnw`) interweaving R code for data generation and \LaTeX code for describing question and solution.
- Exams can be generated by randomly drawing different versions of exercises from a pool of such Sweave exercise templates. The resulting exams can be rendered into various formats including PDF, HTML, **Moodle XML**, or QTI 1.2 (for **OLAT** or **OpenOLAT**).
- Solutions for exercises can be multiple/single-choice answers, numeric values, short text answers, or a combination thereof (cloze).

Exercises

Exercise templates: Sweave files composed of

- R code chunks (within `<>>=` and `@`) for random data generation.
- Question and solution descriptions contained in \LaTeX environments of corresponding names. Both can contain R code chunks again or include data via `\Sexpr{}`.
- Metainformation about type (numeric, multiple choice, ...), correct solution etc. In \LaTeX style but actually commented out.

Simple geometric example:

- Computation of the distance between two points p and q in a Cartesian coordinate system (via the Pythagorean formula).
- Template `dist.Rnw` contained in **exams** package.

```
R> library("exams")
R> exams2pdf("dist.Rnw")
```

Exercises: dist.Rnw

```
<<echo=FALSE, results=hide>>=
p <- c(sample(1:3, 1), sample(1:5, 1))
q <- c(sample(4:5, 1), sample(1:5, 1))
sol <- sqrt(sum((p - q)^2))
@

\begin{question}
What is the distance between the two points
$p = (\$p[1], \$p[2])$ and $q = (\$q[1], \$q[2])$ in a Cartesian coordinate system?
\end{question}

\begin{solution}
The distance $d$ of $p$ and $q$ is given by
$d^2 = (p_1 - q_1)^2 + (p_2 - q_2)^2$ (Pythagorean formula).

Hence $d = \sqrt{(p_1 - q_1)^2 + (p_2 - q_2)^2} = 
\sqrt{(\$p[1] - \$q[1])^2 + (\$p[2] - \$q[2])^2} = 
\$round(sol, digits = 3)$.

[...]
\end{solution}

%% \extype{num}
%% \exsolution{\$round(sol, digits = 3)}
%% \exname{Euclidean distance}
%% \extol{0.01}
```

Exercises: dist.Rnw

```
<<echo=FALSE, results=hide>>=
p <- c(sample(1:3, 1), sample(1:5, 1))
q <- c(sample(4:5, 1), sample(1:5, 1))
sol <- sqrt(sum((p - q)^2))
@

\begin{question}
What is the distance between the two points
$p = (\$p[1], \$p[2])$ and $q = (\$q[1], \$q[2])$ in a Cartesian coordinate system?
\end{question}

\begin{solution}
The distance $d$ of $p$ and $q$ is given by
$d^2 = (p_1 - q_1)^2 + (p_2 - q_2)^2$ (Pythagorean formula).

Hence $d = \sqrt{(p_1 - q_1)^2 + (p_2 - q_2)^2} = \sqrt{(\$p[1] - \$q[1])^2 + (\$p[2] - \$q[2])^2} = \text{\textbackslash round(sol, digits = 3)}$.
[...]
\end{solution}

%% \extype{num}
%% \exsolution{\text{\textbackslash round(sol, digits = 3)}}
%% \exname{Euclidean distance}
%% \extol{0.01}
```

Exercises: dist.Rnw

```
<<echo=FALSE, results=hide>>=
p <- c(sample(1:3, 1), sample(1:5, 1))
q <- c(sample(4:5, 1), sample(1:5, 1))
sol <- sqrt(sum((p - q)^2))
@

\begin{question}
What is the distance between the two points
$p = (\$p[1], \$p[2])$ and $q = (\$q[1], \$q[2])$ in a Cartesian coordinate system?
\end{question}

\begin{solution}
The distance $d$ of $p$ and $q$ is given by
$d^2 = (p_1 - q_1)^2 + (p_2 - q_2)^2$ (Pythagorean formula).

Hence $d = \sqrt{(p_1 - q_1)^2 + (p_2 - q_2)^2} = \sqrt{(\$p[1] - \$q[1])^2 + (\$p[2] - \$q[2])^2} = \text{\textbackslash round(sol, digits = 3)}$.

[...]
\end{solution}

%% \extype{num}
%% \exsolution{\text{\textbackslash round(sol, digits = 3)}}
%% \exname{Euclidean distance}
%% \extol{0.01}
```

Exercises: dist.Rnw

```
<<echo=FALSE, results=hide>>=
p <- c(sample(1:3, 1), sample(1:5, 1))
q <- c(sample(4:5, 1), sample(1:5, 1))
sol <- sqrt(sum((p - q)^2))
@

\begin{question}
What is the distance between the two points
$p = (\$p[1], \$p[2])$ and $q = (\$q[1], \$q[2])$ in a Cartesian coordinate system?
\end{question}

\begin{solution}
The distance $d$ of $p$ and $q$ is given by
$d^2 = (p_1 - q_1)^2 + (p_2 - q_2)^2$ (Pythagorean formula).

Hence $d = \sqrt{(p_1 - q_1)^2 + (p_2 - q_2)^2} = \sqrt{(\$p[1] - \$q[1])^2 + (\$p[2] - \$q[2])^2} = \text{\Sexpr{round(sol, digits = 3)}}$.
[...]
\end{solution}

%% \extype{num}
%% \exsolution{\Sexpr{round(sol, digits = 3)}}
%% \exname{Euclidean distance}
%% \extol{0.01}
```

Exercises: dist.Rnw

```
<<echo=FALSE, results=hide>>=
p <- c(sample(1:3, 1), sample(1:5, 1))
q <- c(sample(4:5, 1), sample(1:5, 1))
sol <- sqrt(sum((p - q)^2))
@

\begin{question}
What is the distance between the two points
$p = (\$p[1], \$p[2])$ and $q = (\$q[1], \$q[2])$ in a Cartesian coordinate system?
\end{question}

\begin{solution}
The distance $d$ of $p$ and $q$ is given by
$d^2 = (p_1 - q_1)^2 + (p_2 - q_2)^2$ (Pythagorean formula).

Hence $d = \sqrt{(p_1 - q_1)^2 + (p_2 - q_2)^2} = \sqrt{(\$p[1] - \$q[1])^2 + (\$p[2] - \$q[2])^2} = \text{\textbackslash round(sol, digits = 3)}$.
[...]
\end{solution}

%% \extype{num}
%% \exsolution{\text{\textbackslash round(sol, digits = 3)}}
%% \exname{Euclidean distance}
%% \extol{0.01}
```

Exercises: L^AT_EX output of Sweave ("dist.Rnw")

```
\begin{question}
What is the distance between the two points
$p = (3, 4)$ and $q = (5, 2)$
in a Cartesian coordinate system?
\end{question}

\begin{solution}
The distance $d$ of $p$ and $q$ is given by
$d^2 = (p_1 - q_1)^2 + (p_2 - q_2)^2$ (Pythagorean formula).

Hence $d = \sqrt{(p_1 - q_1)^2 + (p_2 - q_2)^2} = 
\sqrt{(3 - 5)^2 + (4 - 2)^2}
= 2.828$.

\includegraphics{dist-002}
\end{solution}

%% \extype{num}
%% \exsolution{2.828}
%% \exname{Euclidean distance}
%% \extol{0.01}
```

Exercises: PDF output of exams2pdf ("dist.Rnw")


● Problem

What is the distance between the two points $p = (3, 4)$ and $q = (5, 2)$ in a Cartesian coordinate system?

Solution

The distance d of p and q is given by $d^2 = (p_1 - q_1)^2 + (p_2 - q_2)^2$ (Pythagorean formula).

$$\text{Hence } d = \sqrt{(p_1 - q_1)^2 + (p_2 - q_2)^2} = \sqrt{(3 - 5)^2 + (4 - 2)^2} = 2.828.$$



Exams: Combination of exercises

Idea: An exam is simply a list of exercise templates. For example, using statistics exercise templates contained in **exams**.

```
R> myexam <- list(  
+   "boxplots",  
+   c("confint", "ttest", "tstat"),  
+   c("anova", "regression"),  
+   "scatterplot",  
+   "relfreq"  
+ )
```

Draw random exams:

- First randomly select one exercise from each list element.
- Generate random numbers/input for each selected exercise.
- Combine all exercises in output file(s) (PDF, HTML, ...).

Exams: Combination of exercises

Interfaces: Generate multiple exams via `exams2pdf()`, `exams2html()`, `exams2moodle()`, `exams2qti12()`, ...

Workhorse function: Internally, all interfaces call `xexams()` that handles (temporary) files/directories and carries out four steps.

- ① *Weave*: Each of the selected exercise `.Rnw` files is weaved into a `.tex` file. Default: The standard `Sweave()` function.
- ② *Read*: Each resulting `.tex` file is read into an R list with question, solution, and metainformation. Default: `read_exercise()`.
- ③ *Transform*: Each of these exercise-wise list objects can be transformed, e.g., by converting `LATEX` text to HTML. Default: No transformation.
- ④ *Write*: The (possibly transformed) lists of exercises, read into R for each exam object, can be written out to one or more files per exam in an output directory. Default: No files are written.

Exams: PDF output

`exams2pdf()`:

- The *write* step embeds all questions/solutions into (one or more) master \LaTeX template(s).
- \LaTeX templates control whether solutions are shown, what the title page looks like, etc.
- Compilation of each exam via `pdflatex` (called from within R).

A single exam is popped up in a PDF viewer:

```
R> exams2pdf(myexam, template = "exam")
```

Multiple exams are written to an output directory:

```
R> odir <- tempfile()  
R> set.seed(1090)  
R> exams2pdf(myexam, n = 3, dir = odir,  
+   template = c("exam", "solution"))
```

Exams: PDF output

R University
Statistics Exam 2013-05-18

Exam ID 00001

Name: _____

Student ID: _____

Signature: _____

1. (a) (b) (c) (d) (e)
2.
3.
4. (a) (b) (c) (d) (e)
5. (a) (b) (c) (d) (e)

Statistics Exam: 00001 2

1. In Figure 1 the distributions of a variable given by two samples (A und B) are represented by parallel boxplots. Which of the following statements are correct? (Comment: The statements are either about correct or clearly wrong.)

Figure 1: Parallel boxplots.

(a) The location of both distributions is about the same.
 (b) Both distributions contain no outliers.
 (c) The spread in sample A is clearly bigger than in B.
 (d) The skewness of both samples is similar.
 (e) Distribution A is about symmetric.

2. A machine fills milk into 500ml packages. It is suspected that the machine is not working correctly and that the amount of milk filled differs from the setpoint $\mu_0 = 500$. A sample of 226 packages filled by the machine are collected. The sample mean \bar{y} is equal to 499.7 and the sample variance $s_{\bar{y}-1}^2$ is equal to 576.1.
 Test the hypothesis that the amount filled corresponds on average to the setpoint. What is the absolute value of the t test statistic?

3. For 49 firms the number of employees X and the amount of expenses for continuing education Y (in EUR) were recorded. The statistical summary of the data set is given by:

	Variable X	Variable Y
Mean	58	232
Variance	124	1606

The correlation between X and Y is equal to 0.65.
 Estimate the expected amount of money spent for continuing education by a firm with 60 employees using least squares regression.

4. Figure 2 shows a scatterplot. Which of the following statements are correct?

Exams: HTML output

`exams2html()`:

- In the *transform* step, \LaTeX text is converted to HTML using Ian H. Hutchinson's **TtH** (\TeX to HTML) package.
- Mathematical notation is either represented using MathML (`ttm`), requiring a suitable browser (e.g., Firefox), or plain HTML (`tth`).
- No \LaTeX installation needed, but also limited to \LaTeX commands supported by **TtH**.
- Links to dynamically generated data can be easily included, e.g., `\url{mydata.rda}`.
- The *write* step embeds everything into HTML templates and writes out one HTML file per exam.

A single exam is popped up in a browser, multiple exams are written to an output directory:

```
R> set.seed(1090)
R> exams2html(myexam, n = 3, dir = odir)
```

Exams: HTML output

Screenshot of a web browser window showing a parallel boxplot exercise.

The browser menu bar includes: File, Edit, View, History, Bookmarks, Tools, Help.

The title bar shows "Exam 1".

The address bar shows "file:///tmp/RtmpjJlbQz/file2c8d10dbc901/plain1.html".

The search bar contains "DuckDuckGo".

The main content area displays the following text:

Exam 1

1. Question

In Figure the distributions of a variable given by two samples (A und B) are represented by parallel boxplots. Which of the following statements are correct? (Comment: The statements are either about correct or clearly wrong.)

Figure 1: Parallel boxplots.

Detailed description of Figure 1: The figure is a parallel boxplot with two groups of boxes labeled A and B. Group A consists of two boxes, one for each sample. The left box represents sample A, which has a median at approximately -25.5, an interquartile range (IQR) from about -26.5 to -24.5, and whiskers extending from approximately -20 to -28.5. The right box represents sample B, which has a median at approximately -25, an IQR from about -20 to -25, and whiskers extending from approximately -35 to -15. Both boxes are drawn with black outlines and filled with white space.

Below the plot, there are five statements labeled a through e, each followed by a short description.

- a. The location of both distributions is about the same.
- b. Both distributions contain no outliers.
- c. The spread in sample A is clearly bigger than in B.
- d. The skewness of both samples is similar.
- e. Distribution A is about symmetric.

Exams: Moodle XML

`exams2moodle()`:

- As for HTML output, all \LaTeX text is *transformed* to HTML (plus MathML).
- Rather than writing out one file per exam, a single **Moodle** XML file encompassing all exams is produced.
- All supplementary materials (graphics, data, etc.) are embedded into the HTML code directly using Base64 encoding.
- The resulting `.xml` file can be easily imported into a question bank in **Moodle** and then be used within a **Moodle** quiz.

Multiple replications are written to a single XML file in the output directory:

```
R> set.seed(1090)
R> exams2moodle(myexam, n = 3, dir = odir)
```

Exams: Moodle XML

File Edit View History Bookmarks Tools Help

JSS Quiz

138.232.202.120/mod/quiz/attempt.php?attempt=2 DuckDuckGo

R exams course

Home ► Rexams ► 20 November - 26 November ► JSS Quiz ► Preview

Quiz navigation

1 2 3 4 5

Finish attempt ...

Start a new preview

Navigation


Home My home Site pages My profile Current course Participants Reports General 20 November - 26 November JSS Quiz Info Results 27 November - 3 December 4 December - 10 December 11 December - 17 December 18 December - 24 December 25 December - 31

Question 1

Not yet answered Marked out of 1.00

Flag question Edit question

In Figure 1 the distributions of a variable given by two samples (A and B) are represented by parallel boxplots. Which of the following statements are correct? (Comment: The statements are either about correct or clearly wrong.)



A B

Figure 1: Parallel boxplots.

Select one or more:

a. The location of both distributions is about the same.
 b. Both distributions contain no outliers.
 c. The spread in sample A is clearly bigger than in B.
 d. The skewness of both samples is similar.
 e. Distribution A is about symmetric.

Exams: QTI 1.2 for OLAT

`exams2qti12()`:

- As for HTML output, all \LaTeX text is *transformed* to HTML (plus MathML).
- Rather than writing out one file per exam, a single `.zip` archive is produced, containing the QTI 1.2 XML file plus supplementary materials (graphics, data, etc.) if any.
- Base64 encoding is used for graphics by default, but not for other supplements.
- QTI 1.2 is an international standard for e-learning exams.
- The `.zip` files can be easily imported into **OLAT** (or **OpenOLAT**) when configuring an exam.

Multiple replications are written to a single zipped XML file in the output directory:

```
R> set.seed(1090)
R> exams2qti12(myexam, n = 3, dir = odir)
```

Exams: QTI 1.2 for OLAT

File Edit View History Bookmarks Tools Help

OLAT - OLAT: Course template... +

138.232.202.96:8080/OLAT-LMS-7.6.0.0/auth/1%3A6%3A100020776%3A1% DuckDuckGo Print Help Log out

Home Groups Learning resources Group administration User management Administration gui_demo OLAT: Course...

qt12 Actual score: 0 / 5 Finish test

Still 1 attempt(s).

1.1. Question

1. Exercise
2. Exercise
3. Exercise
4. Exercise
5. Exercise
5.1. Question

Question

In Figure 1 the distributions of a variable given by two samples (A and B) are represented by parallel boxplots. Which of the following statements are correct? (Comment: The statements are either about correct or clearly wrong.)

Figure 1: Parallel boxplots.

a. The location of both distributions is about the same.
b. Both distributions contain no outliers.
c. The spread in sample A is clearly bigger than in B.
d. The skewness of both samples is similar.
e. Distribution A is about symmetric.

Save answer

Exams: QTI 1.2 for OLAT

Caveats: When using exams generated for **OLAT**.

- The text describing the correct solution can only be shown immediately after entering a wrong solution but not after completing the whole exam.
- Numeric exercises are not officially supported by **OLAT**. They do work correctly (with tolerance ranges) but the correct solution is never shown. Hence, by default text matching (with a specific precision and without tolerance ranges) is employed.
- Spaces between columns in matrices have to be enlarged because **OLAT** otherwise collapses them.
- Editing of exercises within **OLAT** does not work.

Discussion

Package exams:

- Framework for automatic generation of simple (mathematical or statistical) exams and associated self-study materials.
- Based on independent exercises in Sweave format which can be compiled into exams (or other collections of exercises).
- Version 1 (Grün and Zeileis 2009) only supported PDF output, version 2 (Zeileis, Umlauf, Leisch 2012) adds an extensible toolbox for various output formats including HTML, **Moodle** XML, and QTI 1.2 (for **OLAT**).
- Contributing to the pool of exercises only requires knowledge of Sweave and minimal markup for metainformation.
- Hosted on R-Forge, providing a support forum:
<http://R-Forge.R-project.org/projects/exams/>

Discussion

At Universität Innsbruck:

- Mathematics course with **OLAT** support (summer/winter term 2012/13 combined: 3,000 participants).
- Team of about 10 persons (professors, lecturers, student assistants) contribute to the pool of exercises.
- During the semester, several online tests (and self tests) are carried out in **OLAT** (via `exams2qti12`) using numerical and multiple-choice exercises.
- Two written exams (via `exams2pdf` with custom template) are carried out using single-choice exercises. Results are scanned by university services and processed by some optical character recognition.
- Instead of generating the PDF files directly, an interface to the “Prüfungsserver” is also available (via `exams2lops`).

References

- Zeileis A, Grün B, Leisch F, Umlauf N (2013). *exams: Automatic Generation of Exams in R*. R package version 1.9-4.
URL <http://CRAN.R-project.org/package=exams>
- Zeileis A, Umlauf N, Leisch F (2012). “Flexible Generation of E-Learning Exams in R: Moodle Quizzes, OLAT Assessments, and Beyond.” *Working Paper 2012-27*, Working Papers in Economics and Statistics, Research Platform Empirical and Experimental Economics, Universität Innsbruck.
URL <http://EconPapers.RePEc.org/RePEc:inn:wpaper:2012-27>.
- Grün B, Zeileis A (2009). “Automatic Generation of Exams in R.” *Journal of Statistical Software*, **29**(10), 1–14.
URL <http://www.jstatsoft.org/v29/i10/>