



Flexible Generation of E-Learning Exams in R: Moodle Quizzes, OLAT Assessments, and Beyond

Achim Zeileis, Nikolaus Umlauf, Friedrich Leisch

<http://eeecon.uibk.ac.at/~zeileis/>

Overview

- Motivation and challenges
- R package **exams**
- Exercises
- Exams
 - Combination of exercises
 - PDF output
 - HTML output
 - XML for **Moodle** or **OLAT**
- Discussion

Motivation and challenges

Motivation:

- Introductory statistics and mathematics courses for business and economics students at WU Wien and Universität Innsbruck.
- Courses are attended by more than 1,000 students per semester.
- Several lecturers teach lectures and tutorials in parallel.
- Need for integrated teaching materials: Presentation slides, collections of exercises, exams, etc.

Challenges:

- *Scalable exams*: Automatic generation of a large number of different exams, both written and online.
- *Associated self-study materials*: Collections of exercises and solutions from the same pool of examples.
- *Joint development*: Development and maintenance of a large pool of exercises in a multi-author and cross-platform setting.

R package exams

Tools chosen: R (for random data generation and computations) and \LaTeX (for mathematical notation) \Rightarrow Sweave.

Design principles of package exams:

- Each exercise template (also called “exercise” for short) is a single Sweave file (`.Rnw`) interweaving R code for data generation and \LaTeX code for describing question and solution.
- Exams can be generated by randomly drawing different versions of exercises from a pool of such Sweave exercise templates. The resulting exams can be rendered into various formats including PDF, HTML, **Moodle XML**, or QTI 1.2 (for **OLAT** or **OpenOLAT**).
- Solutions for exercises can be multiple/single-choice answers, numeric values, short text answers, or a combination thereof (cloze).

Exercises

Exercise templates: Sweave files composed of

- R code chunks (within `<<>>=` and `@`) for random data generation.
- Question and solution descriptions contained in \LaTeX environments of corresponding names. Both can contain R code chunks again or include data via `\Sexpr{}`.
- Metainformation about type (numeric, multiple choice, ...), correct solution etc. In \LaTeX style but actually commented out.

Simple geometric example:

- Computation of the distance between two points p and q in a Cartesian coordinate system (via the Pythagorean formula).
- Template `dist.Rnw` contained in **exams** package.

```
R> library("exams")  
R> exams2pdf("dist.Rnw")
```

Exercises: dist.Rnw

```
<<echo=FALSE, results=hide>>=
p <- c(sample(1:3, 1), sample(1:5, 1))
q <- c(sample(4:5, 1), sample(1:5, 1))
sol <- sqrt(sum((p - q)^2))
@
```

```
\begin{question}
```

What is the distance between the two points

$p = (\text{\Sexpr{p[1]}}, \text{\Sexpr{p[2]}})$ and $q = (\text{\Sexpr{q[1]}}, \text{\Sexpr{q[2]}})$
in a Cartesian coordinate system?

```
\end{question}
```

```
\begin{solution}
```

The distance d of p and q is given by

$d^2 = (p_1 - q_1)^2 + (p_2 - q_2)^2$ (Pythagorean formula).

Hence $d = \sqrt{(p_1 - q_1)^2 + (p_2 - q_2)^2} =$

$\sqrt{(\text{\Sexpr{p[1]}} - \text{\Sexpr{q[1]}})^2 + (\text{\Sexpr{p[2]}} - \text{\Sexpr{q[2]}})^2}$
 $= \text{\Sexpr{round(sol, digits = 3)}}.$

[...]

```
\end{solution}
```

```
%% \extype{num}
```

```
%% \exsolution{\Sexpr{round(sol, digits = 3)}}
```

```
%% \exname{Euclidean distance}
```

```
%% \extol{0.01}
```

Exercises: L^AT_EX output of Sweave("dist.Rnw")

```
\begin{question}
```

What is the distance between the two points

$p = (3, 4)$ and $q = (5, 2)$

in a Cartesian coordinate system?

```
\end{question}
```

```
\begin{solution}
```

The distance d of p and q is given by

$d^2 = (p_1 - q_1)^2 + (p_2 - q_2)^2$ (Pythagorean formula).

Hence $d = \sqrt{(p_1 - q_1)^2 + (p_2 - q_2)^2} =$

$\sqrt{(3 - 5)^2 + (4 - 2)^2}$

$= 2.828$.

```
\includegraphics{dist-002}
```

```
\end{solution}
```

```
%% \extype{num}
```

```
%% \exsolution{2.828}
```

```
%% \exname{Euclidean distance}
```

```
%% \extol{0.01}
```

Exercises: PDF output of exams2pdf ("dist.Rnw")


● Problem

What is the distance between the two points $p = (3, 4)$ and $q = (5, 2)$ in a Cartesian coordinate system?

Solution

The distance d of p and q is given by $d^2 = (p_1 - q_1)^2 + (p_2 - q_2)^2$ (Pythagorean formula).

Hence $d = \sqrt{(p_1 - q_1)^2 + (p_2 - q_2)^2} = \sqrt{(3 - 5)^2 + (4 - 2)^2} = 2.828$.



Exams: Combination of exercises

Idea: An exam is simply a list of exercise templates. For example, using statistics exercise templates contained in **exams**.

```
R> myexam <- list(  
+   "boxplots",  
+   c("confint", "ttest", "tstat"),  
+   c("anova", "regression"),  
+   "scatterplot",  
+   "relfreq"  
+ )
```

Draw random exams:

- First randomly select one exercise from each list element.
- Generate random numbers/input for each selected exercise.
- Combine all exercises in output file(s) (PDF, HTML, ...).

Exams: Combination of exercises

Interfaces: Generate multiple exams via `exams2pdf()`, `exams2html()`, `exams2moodle()`, `exams2qti12()`, ...

Workhorse function: Internally, all interfaces call `xexams()` that handles (temporary) files/directories and carries out four steps.

- 1 *Weave:* Each of the selected exercise `.Rnw` files is weaved into a `.tex` file. Default: The standard `Sweave()` function.
- 2 *Read:* Each resulting `.tex` file is read into an R list with question, solution, and metainformation. Default: `read_exercise()`.
- 3 *Transform:* Each of these exercise-wise list objects can be transformed, e.g., by converting \LaTeX text to HTML. Default: No transformation.
- 4 *Write:* The (possibly transformed) lists of exercises, read into R for each exam object, can be written out to one or more files per exam in an output directory. Default: No files are written.

Exams: PDF output

`exams2pdf()`:

- The *write* step embeds all questions/solutions into (one or more) master \LaTeX template(s).
- \LaTeX templates control whether solutions are shown, what the title page looks like, etc.
- Compilation of each exam via pdf\LaTeX (called from within R).

A single exam is popped up in a PDF viewer:

```
R> exams2pdf(myexam, template = "exam")
```

Multiple exams are written to an output directory:

```
R> odir <- tempfile()
R> set.seed(1090)
R> exams2pdf(myexam, n = 3, dir = odir,
+   template = c("exam", "solution"))
```

Exams: PDF output

R University
Statistics Exam 2013-02-14

Exam ID 00001

Name: _____

Student ID: _____

Signature: _____

1. (a) (b) (c) (d) (e)

2.

3.

4. (a) (b) (c) (d) (e)

5. (a) (b) (c) (d) (e)

Statistics Exam: 00001

2

1. In Figure 1 the distributions of a variable given by two samples (A and B) are represented by parallel boxplots. Which of the following statements are correct? (Comment: The statements are either about correct or clearly wrong.)




Figure 1: Parallel boxplots.

- (a) The location of both distributions is about the same.
(b) Both distributions contain no outliers.
(c) The spread in sample A is clearly bigger than in B.
(d) The skewness of both samples is similar.
(e) Distribution A is about symmetric.
2. A machine fills milk into 500ml packages. It is suspected that the machine is not working correctly and that the amount of milk filled differs from the setpoint $\mu_0 = 500$. A sample of 226 packages filled by the machine are collected. The sample mean \bar{y} is equal to 499.7 and the sample variance s_y^2 is equal to 576.1.
Test the hypothesis that the amount filled corresponds on average to the setpoint. What is the absolute value of the t test statistic?
3. For 49 firms the number of employees X and the amount of expenses for continuing education Y (in EUR) were recorded. The statistical summary of the data set is given by:

	Variable X	Variable Y
Mean	58	232
Variance	124	1606

- The correlation between X and Y is equal to 0.65.
Estimate the expected amount of money spent for continuing education by a firm with 60 employees using least squares regression.
4. Figure 2 shows a scatterplot. Which of the following statements are correct?

Exams: HTML output

`exams2html()`:


- In the *transform* step, \LaTeX text is converted to HTML using Ian H. Hutchinson's **TtH** (\TeX to HTML) package.
- Mathematical notation is either represented using MathML (`ttm`), requiring a suitable browser (e.g., Firefox), or plain HTML (`tth`).
- No \LaTeX installation needed, but also limited to \LaTeX commands supported by **TtH**.
- Links to dynamically generated data can be easily included, e.g., `\url{mydata.rda}`.
- The *write* step embeds everything into HTML templates and writes out one HTML file per exam.

A single exam is popped up in a browser, multiple exams are written to an output directory:

```
R> set.seed(1090)
```

```
R> exams2html(myexam, n = 3, dir = odir)
```

Exams: HTML output



Exam 1

1. Question

In Figure the distributions of a variable given by two samples (A and B) are represented by parallel boxplots. Which of the following statements are correct? (Comment: The statements are either about correct or clearly wrong.)




Figure 1: Parallel boxplots.

- The location of both distributions is about the same.
- Both distributions contain no outliers.
- The spread in sample A is clearly bigger than in B.
- The skewness of both samples is similar.
- Distribution A is about symmetric.

Exams: Moodle XML

`exams2moodle()` :

- As for HTML output, all \LaTeX text is *transformed* to HTML (plus MathML).
- Rather than writing out one file per exam, a single **Moodle** XML file encompassing all exams is produced.
- All supplementary materials (graphics, data, etc.) are embedded into the HTML code directly using Base64 encoding.
- The resulting `.xml` file can be easily imported into a question bank in **Moodle** and then be used within a **Moodle** quiz.

Multiple replications are written to a single XML file in the output directory:

```
R> set.seed(1090)
```

```
R> exams2moodle(myexam, n = 3, dir = odir)
```

Exams: Moodle XML

File Edit View History Bookmarks Tools Help

JSS Quiz

138.232.202.120/mod/quiz/attempt.php?attempt=2 DuckDuckGo

You are logged in as **Nikolaus Umlauf** (Logout)

R exams course

Home ▶ **Rexams** ▶ 20 November - 26 November ▶ JSS Quiz ▶ Preview

Quiz navigation

1 2 3 4 5

Finish attempt ...

Start a new preview

Navigation

Home

- My home
- Site pages
- My profile
- Current course
 - Rexams**
 - Participants
 - Reports
 - General
 - 20 November - 26 November
 - JSS Quiz**
 - Info
 - Results
 - 27 November - 3 December
 - 4 December - 10 December
 - 11 December - 17 December
 - 18 December - 24 December
 - 25 December - 31 December

Question 1
Not yet answered
Marked out of 1.00
Flag question
Edit question

In Figure the distributions of a variable given by two samples (A und B) are represented by parallel boxplots. Which of the following statements are correct? (Comment: The statements are either about correct or clearly wrong.)




Figure 1: Parallel boxplots.

Select one or more:

- a. The location of both distributions is about the same.
- b. Both distributions contain no outliers.
- c. The spread in sample A is clearly bigger than in B.
- d. The skewness of both samples is similar.
- e. Distribution A is about symmetric.

Exams: QTI 1.2 for OLAT

`exams2qti12()`:

- As for HTML output, all $\text{L}^{\text{A}}\text{T}_{\text{E}}\text{X}$ text is *transformed* to HTML (plus MathML).
- Rather than writing out one file per exam, a single `.zip` archive is produced, containing the QTI 1.2 XML file plus supplementary materials (graphics, data, etc.) if any.
- Base64 encoding is used for graphics by default, but not for other supplements.
- QTI 1.2 is an international standard for e-learning exams.
- The `.zip` files can be easily imported into **OLAT** (or **OpenOLAT**) when configuring an exam.

Multiple replications are written to a single zipped XML file in the output directory:

```
R> set.seed(1090)
```

```
R> exams2qti12(myexam, n = 3, dir = odir)
```

Exams: QTI 1.2 for OLAT

File Edit View History Bookmarks Tools Help

OLAT - OLAT: Course templat... +

138.232.202.96:8080/OLAT-LMS-7.6.0.0/auth/1%3A6%3A1000020776%3A1%: DuckDuckGo

Home Groups Learning resources Group administration User management Administration gui_demos OLAT Course... Print Help Log out

qt12 Finish test

Actual score: 0 / 5

qt12

1. Exercise Still 1 attempt(s)

1.1. Question 0/0

2. Exercise 0/0

2.1. Question 0/0

3. Exercise 0/0

3.1. Question 0/0

4. Exercise 0/0

4.1. Question 0/0

5. Exercise 0/0

5.1. Question 0/0

Question

In Figure the distributions of a variable given by two samples (A and B) are represented by parallel boxplots. Which of the following statements are correct? (Comment: The statements are either about correct or clearly wrong.)

Statistic	Sample A	Sample B
Minimum	-28	-35
Q1	-26	-29
Median	-25	-23
Q3	-24	-20
Maximum	-22	-18

Figure 1: Parallel boxplots.

a. The location of both distributions is about the same.

b. Both distributions contain no outliers.

c. The spread in sample A is clearly bigger than in B.

d. The skewness of both samples is similar.

e. Distribution A is about symmetric.

Save answer

Exams: QTI 1.2 for OLAT

Caveats: When using exams generated for **OLAT**.

- The text describing the correct solution can only be shown immediately after entering a wrong solution but not after completing the whole exam.
- Numeric exercises are not officially supported by **OLAT**. They do work correctly (with tolerance ranges) but the correct solution is never shown. Hence, by default text matching (with a specific precision and without tolerance ranges) is employed.
- Spaces between columns in matrices have to be enlarged because **OLAT** otherwise collapses them.
- Editing of exercises within **OLAT** does not work.

Discussion

Package exams:

- Framework for automatic generation of simple (mathematical or statistical) exams and associated self-study materials.
- Based on independent exercises in Sweave format which can be compiled into exams (or other collections of exercises).
- Version 1 (Grün and Zeileis 2009) only supported PDF output, version 2 (Zeileis, Umlauf, Leisch 2012) adds an extensible toolbox for various output formats including HTML, **Moodle XML**, and QTI 1.2 (for **OLAT**).
- Contributing to the pool of exercises only requires knowledge of Sweave and minimal markup for metainformation.
- Hosted on R-Forge, providing a support forum:
<http://R-Forge.R-project.org/projects/exams/>

Discussion

At Universität Innsbruck:

- Mathematics course with **OLAT** support (for 1,600 participants in winter term 2012/13).
- Team of about 10 persons (professors, lecturers, student assistants) contribute to the pool of exercises.
- During the semester, several online tests (and self tests) are carried out in **OLAT** (via `exams2qt112`) using numerical and multiple-choice exercises.
- Two written exams (via `exams2pdf`) are carried out using single-choice exercises. Results are scanned by university services and processed by some optical character recognition.
- Individual solutions to tests/exams are provided to the students on a web portal (via `exams2html`).

References

Zeileis A, Grün B, Leisch F, Umlauf N (2013). **exams**: *Automatic Generation of Exams in R*. R package version 1.9-3.

URL <http://CRAN.R-project.org/package=exams>

Zeileis A, Umlauf N, Leisch F (2012). “Flexible Generation of E-Learning Exams in R: Moodle Quizzes, OLAT Assessments, and Beyond.” *Working Paper 2012-27*, Working Papers in Economics and Statistics, Research Platform Empirical and Experimental Economics, Universität Innsbruck.

URL <http://EconPapers.RePEc.org/RePEc:inn:wpaper:2012-27>.

Grün B, Zeileis A (2009). “Automatic Generation of Exams in R.” *Journal of Statistical Software*, **29**(10), 1–14.

URL <http://www.jstatsoft.org/v29/i10/>